

Effect of magnetic fields on bound localized electron states

F.Kh. Chibirova*

Karpov Institute of Physical Chemistry , 103064, Moscow, Russia

V.R. Khalilov†

Physics Department, Moscow State University, 119899, Moscow, Russia

An approximate analytic solution for the ground electron state are found to the Schrödinger equation for a combination of a uniform magnetic field and single attractive $-\lambda\delta(\mathbf{r})$ potential. Effect of the magnetic field on this bound localized electron state is discussed. We show that above effect leads to appearing the probability current density in some region centered in the point $\mathbf{r} = 0$ as well as to enlarging (for a number of physical models) the localization region of the electron in the plane perpendicular to the magnetic field. We are expected that these properties can be of importance for real quantum mechanical systems.

I. INTRODUCTION

When an electron travels in a uniform magnetic field (that is in a cylindrical external field configuration) where a natural assumption is that the relevant quantum mechanical system is invariant along the symmetry (z) axis, the system then becomes essentially two-dimensional in the xy plane [1]. Many physical phenomena occurring in quantum systems of electrically charged fermions, which have the axial symmetry, can be studied effectively by means of the corresponding Schrödinger or Pauli equation in 2+1 dimensions.

A permanent interest in this topic also is stimulated by the studies of (2+1)-dimensional models in both superconductivity [2] and particle theory (including the quantum Hall effect [3] and the degenerate planar semiconductors with low-energy electron dynamics [4])[2-4].

Two-dimensional electrons in external electromagnetic fields at low temperature has attracted significant interests since the discovery of the integral quantum Hall effect (IQHE) [3]. The amazing fact that the quantized values of the Hall conductance are related only to fundamental constants and are independent of the actual sample and geometry. Another important fact that the IQHE occurs in impure samples with defect centers when localized electron states appear. There are defects of such type in thin films too. It was shown in [5] that the residual effect was observed in the Mössbauer spectra of defect oxide thin films after magnetic field treatment.

The purpose of this paper is to investigate the effect of the magnetic field on planar localized electron states.

II. ELECTRON IN A UNIFORM MAGNETIC FIELD

At first let us consider an electron of charge $e < 0$ in the xy plane in a uniform magnetic field $\mathbf{B}$, which is specified in Cartesian coordinates as

$$\mathbf{B} = (0, 0, B) = \nabla \times \mathbf{A}, \quad \mathbf{A} = (-yB, 0, 0). \quad (1)$$

The Pauli equation in field (1) has the form

$$i\hbar \frac{\partial}{\partial t} \psi(t, \mathbf{r}) = \mathcal{H} \psi(t, \mathbf{r}), \quad \mathbf{r} = (x, y), \quad (2)$$

where the Hamiltonian $\mathcal{H}$ is

$$\mathcal{H} = \frac{1}{2m} \left(-i\hbar \frac{\partial}{\partial x} + \frac{eB}{c} y \right)^2 - \frac{\hbar^2}{2m} \frac{\partial^2}{\partial y^2} + \mu \sigma_3 B. \quad (3)$$

*Electronic address: chibir@cc.nifhi.ac.ru

†Electronic address: khalilov@thc.phys.msu.su

Here $m > 0$ is the effective electron mass in a crystal lattice, $\mu = |e|\hbar/2m_e c$ is the Bohr magneton, m_e is the mass of a free electron and

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The last term in (3) describes the interaction of the spin magnetic moment of the electron with the magnetic field. If the electron is free we shall put $m = m_e$.

Electron wave function in field (1) has the form

$$\psi_{np}(t, \mathbf{r}) = \frac{1}{2} e^{-iE_{ns}t/\hbar} e^{ipx/\hbar} U_n(Y) \begin{pmatrix} 1+s \\ 1-s \end{pmatrix} \quad (4)$$

where

$$E_{ns} = \hbar\omega \left(n + \frac{1}{2} \right) + s\hbar\omega \frac{m}{2m_e} \quad (5)$$

is the energy spectrum of two-dimensional electron, $\omega = |eB|/mc$, $s = \pm 1$ is conserving spin quantum number, p is the momentum of electron in the x -direction in field (1). Note that p is constrained by $|p| \leq eBL/c$ [6].

The functions

$$U_n(Y) = \frac{1}{(2^n n! \pi^{1/2} a)^{1/2}} \exp\left(-\frac{(y-y_0)^2}{2a^2}\right) H_n\left(\frac{y-y_0}{a}\right),$$

are expressed through the Hermite polynomials $H_n(z)$ is the Hermite polynomial, the integer $n = 0, 1, 2, \dots$ indicates the Landau level number, $a = \sqrt{\hbar c/|eB|} \equiv \sqrt{\hbar/m\omega}$ is the so-called magnetic length and $y_0 = -cp/eB$. It should be reminded that the classical trajectory of an electron in the xy plane perpendicular to the magnetic field is a circle with the rest center. Quantity y_0 corresponds to the classical y -coordinate of the circle center.

All electron states (4) are not localized in the x -direction.

III. ELECTRON IN AN ATTRACTIVE POTENTIAL IN THE PRESENCE OF A UNIFORM MAGNETIC FIELD

Now we study a simple solvable model. We consider the motion of an electron in a single attractive $-\lambda\delta(\mathbf{r})$ potential in the presence of a uniform magnetic field. Here λ is a positive constant, $\delta(\mathbf{r})$ is the Dirac delta function. In fact, the equation we need solve is the following Schrödinger equation

$$\frac{1}{2m} \left[\left(-i\hbar \frac{\partial}{\partial x} + \frac{eB}{c} y \right)^2 - \hbar^2 \frac{\partial^2}{\partial y^2} - \hbar^2 \lambda \delta(\mathbf{r}) \right] \Psi_E(\mathbf{r}) = E \Psi_E(\mathbf{r}). \quad (6)$$

Indeed, because the electron spin is conserved, we cannot take into account the last term in (3). Solutions of Eq.(6) are sought in the form

$$\Psi_E(\mathbf{r}) = \sum_{n=0}^{\infty} \int dp C_{Enp} \psi_{np}(\mathbf{r}) \equiv \sum_{n,p} C_{Enp} \psi_{np}(\mathbf{r}), \quad (7)$$

where $\psi_{np}(\mathbf{r})$ is the spatial part of the wave functions (4).

Then for coefficients C_{Enp} , it is easily to obtain

$$C_{Enp}(n+b) = \lambda_0 \sum_{l,k} C_{Elk} V_l(0) V_n(0), \quad (8)$$

where

$$b = \frac{1}{2} - \frac{E}{\hbar\omega}, \quad \lambda_0 = \frac{\lambda}{4\pi m\omega a}, \quad V_l(0) \equiv \sqrt{a} U_l(y=0).$$

Supposing

$$C_{Enp} = C_E \frac{V_n(0)}{n+b}, \quad (9)$$

then inserting (9) in (8) and taking account of formulas

$$\sum_{n,p} |C_{Enp}|^2 = 1, \quad (10)$$

$$\sum_p V_n(0)V_k(0) \equiv \int dp V_n(0)V_k(0) = \frac{\hbar}{a} \delta_{n,k}, \quad (11)$$

we obtain the following equations

$$1 = \frac{\lambda}{4\pi} \sum_{n=0}^{\infty} \frac{1}{n+b}, \quad (12)$$

$$(C_E)^{-2} = \sum_{n=0}^{\infty} \frac{1}{(n+b)^2}. \quad (13)$$

Eq.(12) defines implicitly the energy of a bound localized electron state.

It is seen from Eq. (12) that the electron energy in the lowest unbound state in the presence of uniform magnetic field is equal to $\hbar\omega/2$ and the latter keep on to have the same value in three-dimensional space. Hence, one can conclude that there always exists a bound electron state in any attractive potential in the presence of magnetic field in three-dimensional space. The values of energies certainly differ from each other for two and three spatial dimensions, nevertheless the motion patterns in the plane perpendicular to the magnetic field will be like.

It is pleasant feature of two-dimensional model discussed to make it possible to find both the energy and the approximate wave function of bound state in clear simple forms. At first it is well to note that λ is a dimensionless constant in two-dimensional case so, in fact, Eq.(6) does not contain a term of the energy dimension. Nevertheless, a bound electron state exists in the attractive $-\lambda\delta(\mathbf{r})$ potential. One can see that the sum on the right of Eq.(12) diverges. Introducing the cut-off at $n = N$, we obtain

$$1 = \frac{\lambda}{4\pi} \ln \left(\frac{N}{b} + 1 \right). \quad (14)$$

Thus, we have at large N

$$-E + \frac{\hbar\omega}{2} = \hbar\omega N \exp \left(-\frac{4\pi}{\lambda} \right). \quad (15)$$

In order to consider λ as the bare coupling constant we must require that it should depend on N so as to quantity $-E + \hbar\omega/2$ would remain finite at $N \rightarrow \infty$. Thus, the cut-off parameter N , which tends to infinity, transmutes in arbitrary binding energy $|E|$. This is a nonrelativistic analog of the dimensional transmutation phenomenon which occurs in massless relativistic field theories [7]. For the model under discussion in the absence of magnetic field this phenomenon was first considered in [8].

Now omitting simple calculations we give the final expression for the normalized wave function of bound electron (see, also [9, 10])

$$\Psi_0(\mathbf{r}) = \frac{1}{\sqrt{2\pi}a} \exp \left(\frac{-x^2 - y^2 + 2ixy}{4a^2} \right). \quad (16)$$

In the plane perpendicular to the magnetic field electron is localized in the region $x, y \sim a$. This electron state has a wonderful property. Its probability current densities are

$$j_x = -\frac{\hbar}{2\pi m a^4} y \exp \left(\frac{-x^2 - y^2}{2a^2} \right) \equiv -j_0 y \exp \left(\frac{-x^2 - y^2}{2a^2} \right), \quad j_y = j_0 x \exp \left(\frac{-x^2 - y^2}{2a^2} \right). \quad (17)$$

It follows from Eq. (17) that there is the probability current in the region $x, y \sim a$ but its divergence is equal to zero everywhere. Note that $|\Psi_0(\mathbf{r})|^2$ and two-dimensional current (17) satisfy the continuity equation

$$\frac{\partial |\Psi_0(\mathbf{r})|^2}{\partial t} + \frac{\partial j_x}{\partial x} + \frac{\partial j_y}{\partial y} = 0 \quad (18)$$

everywhere. It follows from here that $|\Psi_0(\mathbf{r})|^2$ is conserved in time. Thus, the state (16) carries but not creates electrical charge. In a definite sense one can say that the localized electron becomes “movable” in the presence of magnetic field.

Results can be treated quasi-classically. Let two-dimensional vector field $\mathbf{j}$ be a complex quantity $\mathbf{j} = j_x + ij_y$ in which components j_x and j_y are functions of the complex variable $z = x + iy$. Note that $\mathbf{j}$ is a solenoidal vector. Computing $\mathbf{c} = [\nabla \times \mathbf{j}]$, we obtain

$$[\nabla \times \mathbf{j}] = \frac{\partial j_y}{\partial x} - \frac{\partial j_x}{\partial y} = \frac{A}{\pi} \left(1 - \frac{zz^*}{d^2} \right) \exp \left(-\frac{zz^*}{d^2} \right), \quad (19)$$

where $A = e\hbar/ma^4$, $d^2 = 2a^2$ and z^* is the complex conjugate of z . The field point in which $\mathbf{c} \neq 0$ is the vortex point or the field vortex. One can write $\mathbf{j}$ in the form

$$\mathbf{j} = \frac{iI(z)}{2\pi z^*}, \quad (20)$$

where

$$I(z) = Azz^* \exp \left(-\frac{zz^*}{d^2} \right) \quad (21)$$

is the vortex intensity that is the vector field circulation on a closed contour encircling the vortex located in the point $x, y = 0$ (see, Figure 1).




FIG. 1: Probability current

It should be borne in mind that for three-dimensional fields $I(z)$ defines only the vortex projection on the normal direction to the plane $z = 0$, while the vortex is three-dimensional vector. For planar fields the vortex can be considered as a vector perpendicular to the plane $z = 0$.

If there are s attractive δ -potentials located in points x_k, y_k then the current vector is equal to the sum

$$\mathbf{j} = \frac{i}{2\pi} \sum_{k=1}^s \frac{I(z_k)}{z - z_k}. \quad (22)$$

It is interest to compare function (16) with the analogous wave function of an electron in a single attractive $-\lambda\delta(\mathbf{r})$ potential in the absence of magnetic field. The latter has been easily found in the form

$$\Psi(\mathbf{r}) = \frac{\lambda C}{2\pi} K_0 \left(\sqrt{\frac{2m|E_0|(x^2 + y^2)}{\hbar^2}} \right), \quad (23)$$

where C is constant, $K_0(z)$ is the MacDonald function of 0th order, E_0 is the electron energy. Now electron is localized in the region $l_0 \sim \sqrt{\hbar^2/2m|E_0|}$. The probability current densities for state (23) are equal to zero everywhere. Hence, the state (23) does not carry electrical charge.

IV. RESUME

We have shown that the effect of magnetic field on localized electron states leads to appearing the probability current in some region as well as to enlarging (for a number of physical models) the localization region of the electron in the plane perpendicular to the magnetic field.

For the effective mass $m = m_e$ the ratio

$$\frac{a}{l_0} \cong 4 \cdot 10^2 \sqrt{\frac{|E_0|}{B}}, \quad (24)$$

where $|E_0|$ and B should be taken in the electron-volt(s) and kilo-gauss(s), respectively. For typical cases $|E_0| \cong 0.1 \div 0.5$ eV, $B \cong 1 \div 50$ kG. The time, which an electron is in three-dimensional potential well of the width $r_0 \ll l_0$ or $r_0 \ll a$ is related to the one out of the well as $r_0^2 \ln^2(r_0/l_0)/l_0^2$ without or r_0^2/a^2 in the presence of magnetic field.

Using the model under discussion one can see a main result of work [5] concerning the decreasing of the number of defects in thin defect oxide films after the magnetic field treatment.

-
- [1] K. Huang, *Quarks, Leptons, and Gauge Fields*, (World Scientific, Singapore, 1982).
 - [2] F. Wilczek, *Fractional Statistics and Anyon Superconductivity*, (World Scientific, Singapore, 1990).
 - [3] R.E. Prange and S.M. Girvin, eds., *The Quantum Hall Effect* (2nd ed.), (Springer-Verlag, New York, 1990).
 - [4] A.M.J. Schakel and G.W. Semenoff, *Phys. Rev. Lett.*, **66**, 2653 (1991).
 - [5] F.Kh. Chibirova, *Phys. of Solid State*, **43**, 1291 (2001).
 - [6] L.D. Landau, E.M. Lifshitz. *Quantum Mechanics*, 2nd ed. (Pergamon, New York, 1978).
 - [7] S. Coleman, E. Weinberg, *Phys. Rev.* **D7**, 1888 (1973).
 - [8] C. Thorn, *Phys. Rev.* **D6**, 39 (1979).
 - [9] R.E. Prange, *Phys. Rev.* **B23**, 4802 (1981).
 - [10] V.R. Khalilov, *Electrons in Strong Electromagnetic Fields* (Gordon & Breach Sci.Pub., Amsterdam, 1996).